



# Plagiarism Checker X Originality Report

**Similarity Found: 9%**

Date: Minggu, September 06, 2020

Statistics: 322 words Plagiarized / 3731 Total words

Remarks: Low Plagiarism Detected - Your Document needs Optional Improvement.

---

Phytochemicals, Nutritional Value, Antioxidant, and Anticoagulant Activity of *Lactuca sativa* L. Leaves and Stems Abstract The main ingredient in the daily salad dish is the leaves of *Lactuca sativa*, while the stems are usually discarded and not eaten. This study was conducted to compare the stems and leaves of two selected *L. sativa* varieties (*L. sativa* var. *longifolia* L. and *L. sativa* var. *capitata* L.)

related to the preliminary phytochemical investigation, nutrient content, DPPH assay, and in vitro anticoagulant activity by determination of **prothrombin time (PT) and activated partial thromboplastin time (aPTT)**. The findings revealed almost the same phytoconstituents in **the leaves and stems** of each variety, such as terpenoids, flavonoids, coumarins, and others.

Exceptionally, tannins have only been detected in the leaves. Compared to the stems, the leaves of both varieties showed statistically significantly higher levels of raw protein and raw fat. For the DPPH assay, the leaves of both varieties have a statistically significantly **higher antioxidant activity than** the stems.

The **leaves and stem extract of the two selected varieties showed a significant prolongation of PT** ( $P < 0.05$  vs. NS = 12:30 s). **On the other hand, the** aPTT test showed a significant increase in aPTT ( $P < 0.05$  vs. NS = 32:44 s) in the stem extract of both varieties, whereas there was no statistically significant increase in aPTT in the leaves of both varieties.

Both parts of each variety have a diversity of phytochemicals and nutrients. The leaves of both varieties have a **higher antioxidant activity than** the stems. While the stems had higher anticoagulant activity than the leaves. Keywords: **Activated Partial**

**Thromboplastin Time** Anticoagulants DPPH Lactuca sativa Prothrombin Time  
INTRODUCTION The administration of prescribed anticoagulant medicines by patients with cardiovascular diseases is very crucial to minimize the risk of thrombosis formation and to maintain the normal blood flowability (Buttar et al.,

2005). **On the other hand, the** serious side effects of those medicines also cause a danger of bleeding in addition to the difficulty in dose monitoring for patients been under anticoagulant therapy (Cordier & Steenkamp, 2012).

Therefore, the orientation toward the plants as a source of cheap anticoagulant drugs with minimum side effects is the desired objective of anticoagulant therapy. One of the most used up leafy vegetables under family Asteraceae is *Lactuca sativa* (Alavi, 1983; Mughrbi & Auzi, 2020), which is vernacularly known as lettuce or khas in Libya. Popularly, the major consuming part in *L. sativa* is the leaves; whereas, the stems are usually discarded and not eaten.

Most of the researches that have been conducted on *L. sativa* focus only on leaves while the juicy stem has a few documented studies. The leaves of *L. sativa* are considered a valuable source of phenolic compounds (Ribas-Agusti et al., 2011) and sesquiterpene lactones (Michalska et al., 2009). Several biological actions of *L. sativa* leaves have been reported such as antioxidant scavenging activity ( Zdravkovic et al.,

2014), anti-inflammatory property (Araruna & Carlos, 2010), analgesic, antidepressant, and in vivo anticoagulant activities (Ismail & Mirza, 2015; Ismail et al., 2017). Few studies have been carried out on the stems exactly on the stem lettuce or Chinese lettuce (*L. sativa* var. *angustana*). From the stem lettuce (Nie et al., 2017; Nie et al., 2018) have separated a sulfated heteropolysaccharide which posses ses an antioxidant, a -amylase inhibitory, and immunomodulatory activities.

About twenty-one **sesquiterpene lactones were isolated** and characterized from the stem lettuce by (Han et al., 2010), some of these compounds have revealed a significant antioxidant activity. Research is applied to the two parts (leaf and stem) of each variety (*L. sativa* var. *longifolia* and *L. sativa* var. *capitata*) to investigate the phytochemical of methanol extracts.

Also, this **study aims to determine the** nutritional value, in vitro antioxidant activity, and in vitro anticoagulant activity. MATERIALS AND METHODS Plant material collection, identification, and processing Varieties (Romaine lettuce and iceberg lettuce) were collected from different markets in Tripoli, Libya, in November 2017. Samples were identified by Dr. Mohamed N.

Abuhadra, a plant taxonomist at the Department of Botany, Faculty of Science, University of Tripoli. The varieties identified have been *L. sativa* var. *longifolia* L. and *L. sativa* var. *capitata* L. with voucher numbers D1 68107922 and D2 68107921, respectively. The leaves and stems of each variety were separated and washed away by distilled water.

The leaves had been dried in shade for two weeks, but the stems had been dried in the oven at 40°C for 16-18 hours. The dried parts were powdered using an electric grinder and stored in polyethylene sealing bags in the refrigerator at 4°C. Extraction of phytochemicals The powdered leaves of 350 g and the powdered stems of 300 g of both varieties were extracted by cold maceration with a powder ratio of 10 g : 100 ml of solvent (Bhat & Al-Daihan, 2014). The powders were extracted twice, using 99% methanol for four days each time, and then filtered.

The filtrates were concentrated at 40°C under reduced pressure by rotary evaporator. The weights of the dry extracts were calculated and the extracts stored in the refrigerator at 4°C. Preliminary phytochemical investigation The methanolic extracts of each part of both varieties have been subjected for the standard procedures for phytochemical screening such as carbohydrates (Singh et al., 2012; Evans, 2009), saponins (Ngbede et al., 2008), terpenoids, steroids (Mir et al.,

2016), flavonoids (Bhandary et al., 2012), tannins (Edeoga et al., 2005), coumarins (Morsy, 2014), and mucilage (Banu & Cathrine, 2015). Estimation of nutritional value The powder of the leaves and the stems of each variety have been analyzed according to the Association of Official Analytical Chemists (Association of Official Agricultural Chemists, 1990; Association of Official Agricultural Chemists, 2000). All the procedures were repeated in duplicate.

Estimation of total ash Total ash was determined by the primary ignition of 4 g of each sample in a pre-weighed crucible until fully carbonized, and then gradually increased the temperature of the muffle furnace to 550°C overnight until carbon-free ash was obtained. The crucible has been weighed. The percentage of total ash has been calculated as follows: Total ash (%) = (Weight of ash/Weight of sample) x 100 Estimation of moisture content The moisture content was determined by the drying method using the oven.

As much as 5 g of dry powder of sample was placed in previously weighed, dry, and clean petri dish for each sample, and then the dish dried in the oven at 105°C for 3-4 hours until reaching the constant weight. The percent of moisture content was

calculated using the following equation: Moisture content (%) =  $(Wt1 - Wt2 / Wt3) \times 100$  Wt1 = Weight of petri dish and sample before drying Wt 2 = Weight of petri dish and sample after drying Wt 3 = Weight of sample.

Estimation of crude protein The Crude protein was determined using the Kjeldahl method. As much as 1 g of each L. sativa powder was placed in a digestion flask containing 3 g of Kjeldahl catalyst and 12 ml of concentrated H<sub>2</sub>SO<sub>4</sub> and was digested for an hour. The flask was then transferred to the distillation unit with 25 ml of 40% NaOH for ammonia release. The flask content was titrated with 0.1

N HCl in the presence of 25 ml of 4% boric acid with two drops of bromocresol green (BCG) and methyl red 5 : 1. The protein percentage was calculated as follows: Crude protein (%) =  $(\text{Sample volume} - \text{blank volume}) \times N \times 14.007 \times 6.25 \times 100 / \text{Sample weight}$  (mg) N = Normality of standard HCl Estimation of crude fat As much as 100 ml of petroleum ether (40-60°C) was poured in the pre-weighed cup.

About 3 g of each sample was transferred into extraction thimble. The extraction process has proceeded in the automated Soxhlet system by two steps, extraction step an hour followed by rinsing step for 30 minute s. Then the cup was dried in the oven at 105°C and weighed.

The percentage of crude fat was calculated using the following formula: Crude fat (%) =  $(\text{Weight of extracted fat} / \text{Weight of sample}) \times 100$  In vitro analysis of antioxidant activity The quantitative DPPH scavenging activity was determined following Kumarasamy et al. (2007) and Kumar et al. (2014) proto cols little The solution DPPH concentration µg/ml prepared. The stock solutions (10 mg/ml) of dry methanolic extracts of L.

sativa were serially diluted to obtain the following concentrations (5, 2.5, 1.25, 0.625, 0.31, and 0.15 mg/ml). About 1 ml of each diluted solution was mixed with 1 ml of DPPH solution and left for 30 minute s in a dark place for complete reaction. After 30 minute s the UV absorbance was recorded by spectrophotometer at 517 nm. L-ascorbic acid was used as a positive control.

The mixture of 1 ml DPPH and 1 ml methanol was utilized as a negative control. The experiment for each extract was measured in duplicate. The results were expressed as a percentage of radical scavenging activity and were calculated by the formula as described by Najafabad & Jamei (2014). The IC<sub>50</sub> value was calculated from the linear regression equation, which is obtained from the plotted graph of % inhibition of DPPH against the different concentrations of extracts. In vitro assay of anticoagulant activity The study pursues (Cordier et al., 2011; Chen et al.,

2014) protocols for assaying the in vitro anticoagulant action of extracts with suitable amendments. Volunteers recruitment Four healthy volunteers of both genders (ages 25-37 years). They were asked to ensure that they had not any previous history for CVDs or medicines administration such as anticoagulant, antiplatelet, and vitamin supplement, etc. Before blood collection, informed consent has been acquired and signed via all volunteers for their participation in this study.

All volunteers have subjected to twice measurements of routine PT and aPTT tests to ascertain the normality of the collected blood. Volunteers recruitment Four healthy volunteers of both genders (ages 25-37 years). They were asked to ensure that they had not any previous history for CVDs or medicines administration such as anticoagulant, antiplatelet, and vitamin supplement, etc. Before blood collection, informed consent has been acquired and signed via all volunteers for their participation in this study.

All volunteers have subjected to twice measurements of routine PT and aPTT tests to ascertain the normality of the collected blood. Blood collection, plasma preparation, and extracts preparation Blood was collected by venipuncture of each volunteer and collected separately into (3.2%) sodium citrate (9 : 1 v/v, blood : anticoagulant sodium citrate) vacuum tube and mixed well.

Immediately, the collected blood was centrifuged at 4000 rpm for 10 minutes to obtain PPP (platelet-poor plasma). Working solutions for each extract with concentrations (1, 2, and 4  $\mu\text{g}/\mu\text{l}$ ) were prepared in normal saline for biological assay. Anticoagulant activity assay Following manufacturer's two PT aPTT separately to at which step of coagulation pathways were inhibited by *L. sativa* extracts.

heparin  $\mu\text{g} / \mu\text{l}$ ) used a positive control for both parameters and regular saline was used as a negative control. Prothrombin time assay In a fusion tube, 25  $\mu\text{l}$  of normal citrated PPP and 2.5  $\mu\text{l}$  of extract solution was added and incubated at 37 °C for 2-3 minutes. Instantly, the time for clot formation (in seconds) was recorded using a stopwatch after the addition of 100  $\mu\text{l}$  of PT reagent.

Activated partial thromboplastin time assay In a fusion tube, 25  $\mu\text{l}$  of normal citrated PPP and 25  $\mu\text{l}$  of extract solution was added and incubated for 2 minutes at 37°C. Then 25  $\mu\text{l}$  of APTT reagent was added and incubated for 2-3 minutes. By using the stopwatch, the time for clot formation (in seconds) was recorded immediately after the addition of 50  $\mu\text{l}$  of (0.025 M)  $\text{CaCl}_2$  solution.

Statistical analysis The results were expressed as mean  $\pm$  S.D with n = 2 using IBM SPSS

statistics 20. The data were statistically analyzed by one-way ANOVA with Tukey's post hoc test. For the anticoagulant activity experiment, the data were statistically analyzed by one-way repeated measures RM- ANOVA with Bonferroni's post hoc test.

The P-value of less than 0.05 was considered as statistically significant. RESULTS AND DISCUSSION Extraction yield After employing a cold maceration method, the yields of L. sativa (each part of each variety) dried methanolic extracts were presented in Table I. The extraction yields of L.

sativa stems were higher than the leaves for both varieties. Since the L. sativa is freshly consumed and rarely cooked, the application of cold maceration as a method of extraction using methanol may be a more suitable choice for extraction. The high yield of methanolic stem extract may be ascribed to the presence of an abundant quantity of sulfated polysaccharides in the stems as observed by (Nie et al., 2017). Table I. Extraction yield of L. sativa methanolic extracts Variety Part Extraction yield (% w/w) Lactuca sativa var. longifolia Leaves 3.67 Stems 17.35 Lactuca sativa var. capitata Leaves 5.73 Stems 16.26 Preliminary phytochemical investigation The methanolic extracts of L.

sativa varieties leave and stem revealed the presence of various bioactive constituents such as carbohydrates, saponins, terpenoids, steroids, flavonoids, coumarins, and mucilage as tabulated in Table II. However, there was a little dissimilarity between the leaves and the stems, i.e. the tannins have been detected only in the leaves. Table II. Extraction yield of L. sativa methanolic extracts Phtochemical Test Lactuca sativa var.

longifolia Lactuca sativa var. capitata Leaf Stem Leaf Stem Carbohydrates Moscs ++ ++ ++ ++ ++ ++ ++ Saponins Foaming test ++ + ++ + Steroids Lieberman-Burchard test ++ ++ ++ ++ Terpenoids Salkowski tes ++ ++ ++ ++ Flavonoids Alkaline reagent test ++ ++ ++ ++ Tannins Ferric chloride test + - + - Mucilage Absolute alcohol + ++ + ++ Coumarins Fluorescence test + ++ + ++ (-) : Negative result, (+) : Positive result, (++) : Strong positive result Estimation of nutritional value Four components have been measured for the leaves and the stems of both L. sativa varieties and the results are presented in Figure 1.

Concerning the findings of crud protein, the leaves of both varieties implied a significant increase ( $P < 0.05$ ) in the percentage of crude protein in comparison to the stems of each variety. The leaves of L. sativa var. longifolia revealed a superior protein content (23.07%). Crude fat results were very low comparing to other nutrients. The values of the leaves of both varieties were significantly higher ( $P < 0.05$ ) than the stems.

The values of total ash were not significantly different ( $P > 0.05$ ) from each other,

despite the leaves of *L. sativa* var. *capitata* showed a little increase amongst other studied powders. Moisture content results were significantly different ( $P < 0.05$ ) between the leaves and the stems of both varieties. Higher moisture content was observed in the stems of both varieties than the leaves. Figure 1.

The nutrients percentage of dried powder for each part of *L. sativa* varieties. Error bars represent SD ( $n=2$ ) Relatively, the leaves and the stems of *L. sativa* varieties can be assumed as a good source for vegetable protein, which is necessary for body development and function. The leaves of *L. sativa* var. *longifolia* revealed a superior protein content (23.07%), which agrees with the observed result (24.78%) by Fujihara et al.

(2001), who have measured the protein content for other vegetables in addition to *L. sativa* leaves, and the results exhibited that the *L. sativa* has an elevated protein content in relative to *Solanum melongena*, *Daucus carota*, *Brassica rapa*, and *Cucurbita maxima*. Both parts of *L. sativa* are considered as a low-fat diet; therefore, they can be recommended for overweight individuals that are seeking weight loss. The value of *L. sativa* var.

*capitata* leaves showed a greater fat content (3.53%), which is approaching the observation (4.1%) that has been reported by Wheeler et al. (1994). In-vitro analysis of antioxidant activity The efficiency of the sample in scavenging the DPPH radicals is expressed by the IC50 value. A small IC50 value implies a high antioxidant activity. A significant difference ( $P < 0.05$ ) in the IC50 values were found between the leaves and the stems of both varieties. Specifically, the leaves of *L. sativa* var. *capitata* displayed the lowest IC50 (2.41 mg/ml) in comparison to the leaves of *L. sativa* var. *longifolia* (3.23 mg/ml).

However, there was no significant difference in the stems of both varieties with ( $P > 0.05$ ), and they displayed higher IC50 values comparing to the leaves. In comparison to the L- ascorbic acid, the resulted IC50 is (0.01 mg/ml), which is the lowest value among the tested *L. sativa* extracts; therefore, L-ascorbic acid has a significant ( $P < 0.05$ ) superior antioxidant activity more than other tested extracts Figure 2. Figure 2.

IC50 values for each part of *L. sativa* different varieties and L. ascorbic acid. Error bars represent SD ( $n=2$ ) The diversity of *L. sativa* phytoconstituents can be considered as a functional food and adjuvant treatment for several chronic diseases as CVDs. Hence, the antioxidant activity of *L. sativa* may be related to the presence of flavonoids, coumarins, and tannins as many studies have reported the strong correlation between these compounds and the high DPPH scavenging activity of *L. sativa* leaves (Llorach et al.,

2008; Liu et al., 2007).

In this work, a higher antioxidant activity has been observed in leaves in comparison to the stems. This may be attributed to the presence of polyphenolic compounds (tannins) in the leaves. Tannins act as strong natural antioxidants and a substitute for many synthetic antioxidants (Sung et al., 2012).

Under the same conditions of DPPH assay, Gan and Azrina (2016) have measured the IC<sub>50</sub> of 70% aqueous ethanolic extracts of *L. sativa* var. *longifolia* and *L. sativa* var. *capitata* leaves. The resulted values were (4.48 and 3.99 mg/ml), respectively, which are approaching those reported in the present work (3.23 mg/ml and 2.41 mg/ml) for the leaves methanolic extract for the same varieties.

Therefore, the bit dissimilarity in the results may be due to the difference in the solvent of extraction and the phenolic content. In-vitro analysis of anticoagulant activity Coagulation waterfall is mediated by clotting factors, which circulate in blood plasma in an inactive form and are activated in response to any triggers. Coagulation cascade includes two correlating pathways; intrinsic and extrinsic pathways (Triplett, 2000; Palta et al., 2014).

Prothrombin time test reflects the extrinsic pathway because the PT reagent composes of calcified thromboplastin, which stimulates the clotting factors that mediate the extrinsic pathway; whereas, the intrinsic pathway can be examined by aPTT test using aPTT reagent, which is consisted of kaolin cephaline that acts as an activator in presence of calcium ions for intrinsic clotting factors (Jesonbabu et al., 2012).

In this study, the leaves and the stems extract of both selected varieties exhibited a significant prolongation of PT ( $P < 0.05$  vs. NS = 12:30 s). Hence, both the leaves and the extract of the stem of each variety may lead to the reduction coagulation Specifically, post test that concentration has significantly raised the PT ( $P < 0.05$  vs. NS) for both parts of each variety. The leaves extract of *L. sativa* var. *capitata* showed the greatest PT value (27.38 s) among other extracts as presented in Figure 3. Figure 3. Prothrombin time values comparison among the *L. sativa* methanolic extracts with NS and heparin.

Error bars represent the SD (n=4) For the aPTT test, the stems extract of both varieties displayed a significant increase in aPTT ( $P < 0.05$  vs. NS= 32:44 s); whereas, the leaves of both varieties have not exhibited any significant elevation in aPTT at the three concentrations ( $P > 0.05$  vs. NS). Only the concentration 4  $\mu\text{g}/\mu\text{l}$  has significantly prolonged the aPTT ( $P < 0.05$  vs. NS). The stems *L. sativa* var.

longifolia showed a higher aPTT value (43:19 s) than the stems of *L. sativa* var. capitata (40:33 s). The aPTT and PT values of heparin were significantly higher ( $P < 0.05$ ) than other extracts as illustrated in Figures 3 and 4. Figure 4. Activated partial thromboplastin time values comparison among the *L. sativa* methanolic extracts with NS and heparin.

Error bars represent the SD ( $n=4$ ) This research was the first attempt to investigate the in vitro anticoagulant activity of *L. sativa* on human plasma. Sometimes in vitro activity has not been translated into in vivo activity due to the pharmacokinetic considerations. However, Ismail and Mirza (2015) have conducted an in vivo evaluation of the anticoagulant activity of *L. sativa* leaves.

The results displayed a significant anticoagulant activity of *L. sativa* leaves extract. This in vivo study can improve the accuracy of the present study results from the in vitro assay. Hence, we can rely on the in vitro anticoagulant essay as a model for the evaluation of the anticoagulant activity. The presence of various phytoconstituents *L.*

*sativa* extracts such as flavonoids, saponins, terpenoids, steroids, and coumarins, which play an integral role in the reduction of coagulation time (Duric et al., 2015; Ashwini & Asha, 2017; Moghadamtousi et al., 2013). Only the stems of each variety exhibited an effect on the intrinsic pathway by prolongation of aPTT. This may be due to the presence of high content of sulfated polysaccharides in the stems of *L. sativa* relative to the leaves (Nie et al., 2017).

Sulfated polysaccharides have exhibited an effect only on the intrinsic pathway by increasing the aPTT, and they have not displayed any effect on the extrinsic pathway (Rodrigues et al., 2011; Karaki et al., 2013). CONCLUSION From this study, the yield of stem methanol extracts was higher than the yield of leaf for both varieties. Both parts of each variety showed the presence of the same phytochemicals with a slight discrepancy between the leaves and the stems, i.e.

tannins were detected only in the leaves. The content of the leaves' nutrients in both varieties showed statistically significantly higher levels of raw protein, raw fat, and fairly total ash compared to the studied stem powders. As regards the DPPH assay, the leaves of both varieties have a statistically significantly higher antioxidant activity than the stems.

The stems revealed a statistically significant higher anticoagulant activity than the leaves. ACKNOWLEDGMENT Special appreciation to the University of Tripoli and Food Quality Control Center, Tripoli, Libya, for the opportunity to conduct nutritional value experiments. Equally thanks to Mr. Abduladeem Hedwar, Mr. Khaled Smein, Miss. Amina

Osta, and Dr. Abdulrazag Tajuri, for their valuable assisting.

#### INTERNET SOURCES:

---

1% - [www.medscape.com/answers/2085837-179654](http://www.medscape.com/answers/2085837-179654)  
<1% - NBC Sports via Yahoo Sports  
3% - [www.sciencedirect.com/science/article](http://www.sciencedirect.com/science/article)  
<1% - [www.science.gov/topicpages/d](http://www.science.gov/topicpages/d)  
<1% - [www.medicinenet.com/primary\\_biliary\\_cirrhosis](http://www.medicinenet.com/primary_biliary_cirrhosis)  
<1% - [naldc.nal.usda.gov/download/13230](http://naldc.nal.usda.gov/download/13230)  
1% - [www.ncbi.nlm.nih.gov/pmc/articles](http://www.ncbi.nlm.nih.gov/pmc/articles)  
<1% - [www.igntu.ac.in/syllabus/MSc\\_Botany\\_Syllabus](http://www.igntu.ac.in/syllabus/MSc_Botany_Syllabus)  
<1% - [www.wikihow.com/Dry-Rosemary](http://www.wikihow.com/Dry-Rosemary)  
<1% - [shodhganga.inflibnet.ac.in/bitstream/10603/5449/9](http://shodhganga.inflibnet.ac.in/bitstream/10603/5449/9)  
<1% - [ijpsr.com/bft-article/phyto-pharmacognostical](http://ijpsr.com/bft-article/phyto-pharmacognostical)  
<1% - [www.ijcmas.com/vol-2-11/C](http://www.ijcmas.com/vol-2-11/C)  
<1% - [en.wikipedia.org/wiki/AOAC\\_International](http://en.wikipedia.org/wiki/AOAC_International)  
<1% - [link.springer.com/article/10](http://link.springer.com/article/10)  
<1% - [famousminerals.com/testingprocedures](http://famousminerals.com/testingprocedures)  
<1% - [www.sciencedirect.com/engineering/crude-protein](http://www.sciencedirect.com/engineering/crude-protein)  
<1% - [onlinelibrary.wiley.com/doi/full](http://onlinelibrary.wiley.com/doi/full)  
<1% - [www.chegg.com/homework-help/questions-and](http://www.chegg.com/homework-help/questions-and)  
<1% - [www.reddit.com/r/Mcat](http://www.reddit.com/r/Mcat)  
<1% - [www.intechopen.com/online-first/tannins](http://www.intechopen.com/online-first/tannins)  
<1% - [salkowski test flavonoids alkaline reagent test tannins ferric chloride test - - mucilage absolute alcohol coumarins fluorescence](http://salkowski-test-flavonoids-alkaline-reagent-test-tannins-ferric-chloride-test--mucilage-absolute-alcohol-coumarins-fluorescence)  
<1% - [healthyliving.azcentral.com/percentage-fat](http://healthyliving.azcentral.com/percentage-fat)  
<1% - [www.nature.com/articles/srep16851](http://www.nature.com/articles/srep16851)  
<1% - [www.skintrends.com/l-ascorbic-acid](http://www.skintrends.com/l-ascorbic-acid)  
<1% - [firstaidteam.com/2014/11/17/mnemonic-monday](http://firstaidteam.com/2014/11/17/mnemonic-monday)  
<1% - [www.ahajournals.org/doi/10](http://www.ahajournals.org/doi/10)  
<1% - [downloads.hindawi.com/journals/ijfs](http://downloads.hindawi.com/journals/ijfs)  
<1% - [uot.edu.ly/en/university-tripoli-first-libya](http://uot.edu.ly/en/university-tripoli-first-libya)