



Plagiarism Checker X Originality Report

Similarity Found: 11%

Date: Minggu, September 06, 2020

Statistics: 326 words Plagiarized / 3098 Total words

Remarks: Low Plagiarism Detected - Your Document needs Optional Improvement.

Antibiotics Resistance Pattern of Coliform Bacteria Isolated from Slaughterhouse Wastewater in Jega Town, Kebbi State, Nigeria Abstract Wastewater is an essential reservoir of pathogenic bacteria, which include resistant strains. This study determined the antibiotics resistance pattern of Coliform bacteria isolated from slaughterhouse wastewater in Juga local government.

Five different samples of wastewater from different locations of a slaughterhouse were collected using standard sample collection techniques. Coliform bacteria were isolated using the standard microbiological method. The total bacterial count was determined using plate count agar. Suspected coliforms were identified based on morphological and biochemical tests.

Antimicrobial susceptibility testing was carried out using the standard disc diffusion method. The highest bacterial count (8.4×10^2 CFU/ml) was observed from the wastewater collection point, and the lowest bacterial (1.2×10^2 CFU/ml) count from the slaughtering point. Three Gram- negative bacteria, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Enterobacter aerogenes*, were identified.

Escherichia coli was the most frequently isolated in slaughterhouse wastewater 39 (45.8%). Among the antibiotics tested against isolated bacteria, Septrin was the most resistant antibiotics recorded against *E. coli* and *E. aerogenes* with 84.61% and 88.89% resistant, respectively. *Pseudomonas aeruginosa* was found to show higher resistance to Chloramphenicol and Septrin with 84.21% resistance each.

The occurrence of antibiotic-resistant bacteria from slaughterhouse wastewater showed the risks associated with antimicrobial drug resistance transferred from food- producing

animals to humans. Management concerns, such as local government health officers and community development officers, should increase the sensitization of slaughterhouse workers by organizing conferences or conducting radio talk to educate these slaughterhouse workers on the treatment of slaughterhouse wastewater and health risk associated with antimicrobial-drug resistance transferred from animals to humans.

Keywords: Antibiotics Coliform bacteria Jega Resistant Slaughterhouse Wastewater
INTRODUCTION The wastewater is an essential reservoir of bacterial pathogens, including resistant isolates (Egesi et al., 2019). Wastewater from the slaughterhouse, if not carefully treated or disposed of, can be good sources of antibiotics resistant bacteria (Manyi-Loh et al.,

2018), mainly where slaughterhouse is located near the vegetable farms or community sources of drinking water (Egesi et al., 2019). Activities of meat processing from slaughterhouses in Nigeria are mostly handled by people who are unaware of the slaughterhouse and environmental safety policies (Johnson & Etokidem, 2019), especially regarding possible microbial contaminations during meat processing (Egesi et al., 2019).

Wastewater from slaughterhouse may contain a higher level of nutrients such as manure, blood, fats, grease, hair, grit, undigested feeds, salts, phosphates, and nitrates support the growth of microbes, including bacterial pathogens (Kundu et al., 2013; Egesi et al., 2019). Bacteria pathogens from animal sources, especially coliform, acquire by a human was reported to be more challenging to treat (Hiroi et al., 2012).

The reason is that current technology in animal food production involved antimicrobial agents in enhancing the production (Hiroi et al., 2012; Egwuenu et al., 2018). The most commonly isolated bacteria from slaughterhouse wastewater are *Salmonella* spp., *Escherichia coli* O157 : H7, *Shigella* spp., *Staphylococcus aureus*, *Streptococcus* spp., *Yersinia enterocolitica*, and *Campylobacter* spp. (Nafarnda et al., 2012). Similarly, Adebowale et al.

(2016) reported the occurrence of *Enterobacter aerogenes*, *Hafnia alvei*, *Erwinia mallotivora*, *Edwardsiella ictaluri*, *Enterobacter amnigenus*, *E. coli* O157 strains, *Proteus mirabilis*, *Staphylococcus* spp., *Pseudomonas aeruginosa*, *Enterobacter intermedius*, *Yersinia aleksiciae*, *Serratia odorifera*, and *Enterobacter cloacae* from the central slaughterhouse and its receiving surface water in Abeokuta, Nigeria. Ogbomida et al. (2016) also reported *Escherichia* sp., *Pseudomonas* sp., *Enterobacter* sp., *Klebsiella* sp., *Staphylococcus* sp.,

Salmonella sp., and Streptococcus sp. from slaughterhouse wastewater. Literature has shown that some of these bacterial species were recently classified as ESKAPE (Enterococcus faecium, Staphylococcus aureus, Klebsiella pneumoniae, Acinetobacter baumannii, Pseudomonas aeruginosa, and Enterobacter spp.)

bacterial pathogens (Santajit & Indrawattana, 2016; World Health Organization, 2017; Shrivastava et al., 2018; Savin et al., 2020). In Nigeria, monitoring environmental sanitation and waste management were established and implemented (Egwuenu et al., 2018). However, the agency lacks monitoring to assure the proper practice of these policies by some local community, which leads to the spread of these commonly resistant bacteria within the community.

Research on resistant bacteria is fundamental as this will provide some preliminary information about the existence of these bacteria within the local community (Aslam et al., 2018). However, literature has shown a lack of comprehensive data regarding resistance to antimicrobial agents, especially in African continents, including Nigeria (Egwuenu et al., 2018). Therefore, this study was designed to determine the antibiotics resistance pattern of coliform bacteria isolated from slaughterhouse wastewater.

MATERIALS AND METHODS Research sites This study was carried out in the Jega local government area of Kebbi State (Figure 1). The local government is located at the latitude and longitude 12.3667°N 4.6333°E. The inhabitants are predominantly Hausa people by the tribe. The primary occupation of the people is farming and trading. The local government has an area of 891 km² and a population of 193,352 at the 2006 census.

Jega local government has a local market that traditionally holds on Friday, but business activities even the The had animal's point, a processing meat point where neighboring local governments buy and sell animals and meats, especially during market day. The market serves as sources of meats for many neighboring local government areas — the people of Jega local government slaughterhouse butchering cattle, goats, sheep, and camel. Sample collection A total of five samples were collected from different slaughterhouse sampling points.

Samples were collected using the grab sampling method with a wide-mouthed 250 mL sterilized Pyrex glass bottles with tight screw dustproof stoppers. Slaughterhouse wastewater samples were placed in a cooler box and transported to the microbiology laboratory at the Department of Microbiology, Faculty of Life Sciences, Kebbi State University of Science and Technology, Aliero for further analysis. Figure 1. Map of Kebbi

State Nigeria, showing the research sites (Ajibola et al.,

2018) Isolation and identification of coliform bacteria Coliform bacteria were isolated using the surface spread plate method described by Cheesbrough (2006) and Bashir et al. (2018). Total coliforms and bacterial counts of the collected wastewater samples were determined using Eosin Methylene Blue (EMB) agar and Plate count agar.

Serial dilutions of the water samples were made aseptically by transferring 1 mL of the wastewater sample into 9 mL of sterile distilled water and diluting that range from 10^{-1} to 10^{-6} . After that, an aliquot of 0.1 mL from 10² dilutions was plated on Plate count agar (Hi-media) in duplicates then spread aseptically using a sterile spreader (Egesi et al., 2019).

Plates were incubated at 37°C for 24 hours and colonies were counted and expressed as colonies forming units per milliliter (CFU/mL). Coliform counts of wastewater samples were determined using the three-tube analysis of Most Probable Number (MPN) techniques, as described earlier by Cheesbrough (2006) and Bashir et al. (2018) using sterile MacConkey broth.

The first set of three tubes had sterile 10 mL double-strength lactose broth, whereas the second and third sets had 10 mL single strength broth. All tubes contained Durham tubes before sterilization. The three sets of tubes received 1 mL, 0.1 mL, and 0.01 mL quantities of wastewater samples. The experiment was conducted in duplicates.

Tubes were then incubated at 37°C for 24-48 hours to estimate total coliforms and, after that, were examined for acid and gas production. Acid production was detected by the color change of the lactose broth from reddish. A loopful of samples from positive tubes were inoculated onto Petri plates containing Eosin Methylene blue (EMB) agar using the streaking method, and then plates were incubated at 37°C for 24 - 28 hours. This selective and differential agar medium inhibits Gram-positive bacteria but allows the growth of Gram-negative coliforms.

Coliforms produced large pinkish colonies, with dark centers and green metallic shine. This confirmative test was carried out to determine if the coliforms were of the fecal origin or not. The presence of typical colonies at 37°C confirmed coliforms, as referenced by Cheesbrough (2006) and Bashir et al. (2018).

The suspected coliform isolates were further confirmed using convention biochemical tests such as the citrate test, indole test, Urease test, Methyl red test, Voges Proskauer test, Triple sugar iron test. All identification processes were carried out with the

guidance from [Bergey's Manual of Determinative Bacteriology](#) (Holt et al., 1994).

Antimicrobial susceptibility test Antibiotic susceptibility patterns of the pure colony of *E. coli*, *P. aeruginosa*, and *E. aerogenes* isolates were determined using the disc diffusion method. The antibiotic discs used in this study were Ciprofloxacin (10 µg), Gentamycin (10 µg), Augmentin (30 µg), Pefloxacin (20 µg), Amoxicillin (10 µg), Streptomycin (30 µg), Sparfloxacin (10 µg), Ofloxacin (30 µg), Septrin (30 µg), and Chloramphenicol (30 µg). Inoculum suspension of a test organism (*E. coli*, *P.*

aeruginosa, and *E. aerogenes*) was prepared by inoculating pure colonies into sterile test tubes containing 2 mL of normal saline. The suspension was then adjusted by standardizing the turbidity to that of 0.5 McFarland standards. The standardized suspension was inoculated on the sterile plates containing freshly prepared Mueller Hinton agar using a sterile cotton swab.

The surfaces of the plates were then impregnated with the antibiotic discs using a sterile Forceps. The plates were incubated for 24 hours at 37°C. After incubation, the inhibition zones were measured in mm using a ruler by the Clinical and Laboratory Standards Institute (2019). RESULTS AND DISCUSSION A study on antimicrobial-resistant is critical now a day mainly from the African continent, where comprehensive data to show the magnitude of resistance are lacking (Egwuenu et al., 2018).

Literature has shown that wastewater from slaughterhouse can be served as a source or reservoir of resistant bacterial pathogens spread within the community if not well treated or disposed of (Egesi et al., 2019). This resistance varies from different locality due to some factors such as the educational background of animal rearers, use of antibiotics in animal feedstuffs, among others (Egwuenu et al., 2018).

This hinders the effort to control antimicrobial-resistant humans because resistant bacteria of animal origin are more dangerous than human origin (Hiroi et al., 2012). This research determined the antibiotics resistance pattern of coliform bacteria isolated from slaughterhouse wastewater in Jega local government area Kebbi State, Nigeria.

Out of five samples of wastewater collected from different slaughterhouse points, the result showed that the wastewater collection point had the highest bacterial coliform counts with a mean and standard deviation of $(8.73 \pm 0.35 \times 10^2 \text{ CFU/mL})$ while the Slaughtering point had the lowest coliform bacterial counts with mean and standard deviation of $(1.27 \pm 0.31 \times 10^2 \text{ CFU/mL})$ as presented in Table I. This was in line with the study of Jega et al.

(2018), who reported coliform bacteria from slaughterhouse Effluent-Impacted Tagangu River, Aliero, Kebbi State, North- Western Nigeria. The bacterial coliform count reported in this study was similar to the findings of Atuanya et al. (2018) who reported mean and standard deviation of bacterial coliform count ranged $2.7 \times 10^5 \pm 0.1$ to $3.8 \times 10^5 \pm 1.4$ CFU/mL and $5.5 \times 10^5 \pm 0.3$ to $9.2 \times 10^5 \pm 0.4$

CFU/mL from government and private slaughterhouse from Effluents around Ikpoba River in Benin City, Nigeria. However, Egesi et al. (2019) also reported bacterial coliform from different slaughterhouse points. Table I. The mean and standard coliform bacterial count of different slaughterhouse wastewater sample in Jega government area, Kebbi States, Nigeria Slaughterhouse points Mean \pm SD coliform counts ($\times 10^2$ CFU/mL) p-value Wastewater collection point 8.73 ± 0.35 0.03671 Washing point 1.77 ± 0.46 Drainage 1.87 ± 0.44 Slaughtering point 1.27 ± 0.31 Un-skinning point 1.3 ± 0.2

CFU: Colony-forming unit This research's findings showed higher coliform count from the wastewater collection point compares to either point of the slaughterhouse. The comparison between the different slaughterhouse points using Tukey's multiple comparisons showed statistically different ($p = 0.03671$). This finding is similar to Egesi et al. (2019), who reported higher coliform bacterial counts from slaughterhouse effluents than slaughterhouse sampling points.

The suspected coliform bacteria were identified using a standard biochemical test, as shown in Table II. Among the **coliform bacteria isolated from** the different slaughterhouse sampling points, E. coli is the most abundant bacteria isolated with 39 points (45.8%), as presented in Table III. The presence of E.

coli from slaughterhouse wastewater is an indication of fecal contamination, and therefore such water needs to be given special treatment to arrest the spread of this pathogenic organism within the environment. Disease caused by E. coli of animal origin is more difficult to treat, especially the resistant isolate due to the frequent use of antibiotics in animal feeds (Hiroi et al., 2012).

This **is in line with the** finding of Jega et al. (2018) who reported the E. coli as the most abundant **bacteria isolated from slaughterhouse** effluent-impacted Tagangu River, Aliero, Kebbi State, North-Western Nigeria. Similarly, Egesi et al. (2019) reported E.

coli as the second most abundant **coliform bacteria isolated from slaughterhouse effluents** discharged into water bodies in Owerri, Nigeria. However, Tesfaye et al. (2019) reported E. coli as the most abundant coliform bacteria from **wastewater samples from health facilities**, slaughterhouses, downstream rivers, and a wastewater treatment plant

in Addis Ababa, Ethiopia. Other Gram-negative bacteria isolated from slaughterhouse wastewater are *P. aeruginosa* and *E. aerogenes*. The presence of *P.*

aeruginosa in slaughterhouse wastewater was also reported by previous studies (Igbinosa et al., 2012; de Oliveira et al., 2013; Ayogu et al., 2018). This confirmed the statement by Yayan et al. (2015) that *P. aeruginosa* is currently one of the dangerous nosocomial organisms that can be transmitted in many ways, including food products such as meat. The percentage (22.4%) of *P. aeruginosa* reported in this study is lower than the 37.6% reported by Rabiun and Falodun (2017) from wastewater of a slaughterhouse in Ibadan, Nigeria.

Table II. Biochemical characteristics of coliform bacteria isolated from slaughterhouse waste-water in Jega local government area, Kebbi States Nigeria

Parameters	Strains results 1	2	3
Gram's staining	(rod)	(rod)	(rod)
Citrate	+	-	+
Motility	+	+	-
Oxidase	-	+	+
Catalase	+	+	+
Indole	-	+	-
Urease	-	-	-
Methyl red	+	+	-
Voges Proskauer	+	-	-
Triple sugar iron Slant	Alkaline	Acid	Alkaline
But Acid	Acid	Acid	Acid
H ₂ S	-	-	-
Starch	-	+	-
Gelatin	+	-	+
Sugar fermentation	Glucose + / Acid	+ / Acid	- Maltose + / Acid + / Acid -
Sucrose	+ / Acid	+ / Acid	- Lactose - + / Acid -
Mannitol	+ / Acid	+ / Acid	+ / Acid
Possible microorganism	<i>E. aerogenes</i>	<i>E. coli</i>	<i>P.</i>

aeruginosa Table III. Coliform bacteria isolated from wastewater samples from different slaughterhouse points in Jega local government area, Kebbi States Nigeria

Microorganism	Slaughterhouse points n (%)	Washing point n (%)	Drainage n (%)	Slaughtering point n (%)	Un - skinning point n (%)	Total n (%)	p value	Wastewater collection point
<i>E. coli</i>	18 (46.2)	8 (20.5)	1 (2.6)	5 (12.8)	7 (17.9)	39 (45.8)	0.5057	
<i>E. aerogenes</i>	14 (51.9)	2 (7.4)	5 (18.5)	6 (22.2)	0 (0)	27 (31.8)		
<i>P. aeruginosa</i>	10 (52.6)	0 (0)	3 (15.8)	1 (5.3)	5 (26.3)	19 (22.4)		
Total	42 (49.4)	10 (11.8)	9 (10.6)	12 (14.1)	12 (14.1)	85 (100)		

The result of antibiotics resistant pattern of the coliform bacteria isolated (Figure 2) indicates that *E.*

coli showed higher resistance to Septrin and Augmentin among the antibiotics tested with (84.61%) and (89.74%) resistant, respectively. The higher resistance of *E. coli* recorded in this research is worrying as recent literature showed that many of the antimicrobial-resistant strains of *E. coli* that cause human urinary tract and bloodstream infections appear to have arisen from contaminated meat products (Egwuenu et al., 2018).

The finding of this research is in line with the report of Atuanya et al. (2018) and Savin et al. (2020), who reported the resistance of *E. coli* isolated from slaughterhouse Effluents to Septrin and Augmentin. Similarly, Jega et al. (2018) also reported the resistance of *E. coli* isolated from slaughterhouse Effluent-Impacted Tagangu River, Aliero, to

Augmentin. However, Tesfaye et al. (2019) reported the resistant E.

coli isolated from e in wastewater samples from health facilities, slaughterhouse, downstream rivers, and a wastewater treatment plant in Addis Ababa, Ethiopia to Septrin including multi-drugs resistant isolate. Enterobacter aerogenes showed resistance to Septrin (88.89%). Enterobacter aerogenes were reported to be an opportunistic nosocomial pathogen and are sought out to be one of the many critical causes for extra-intestinal infections next to E. coli.

The infection associated with this organism is respiratory and urinary tract infections, specifically cystitis, and wound, bloodstream, and central nervous system infections (Flores-Mireles et al., 2015). The finding of this research is in line with the report of Jega et al. (2018), who also reported the resistance of Enterobacter spp. isolated from slaughterhouse Effluent-Impacted Tagangu River, Aliero to Augmentin and Septrin.

The results of this finding showed that P. aeruginosa is highly resistant to Chloramphenicol and Septrin with 84.21% resistant each. These results are in line with Rabiun and Falodun's (2017) finding, who reported 40.6% resistance of P. aeruginosa isolated from wastewater of a slaughterhouse in Ibadan, Nigeria, to Chloramphenicol.

This bacterium's resistance to this drug is alarming as this bacterium was reported to be one of the dangerous nosocomial microorganisms that can be transmitted in many ways, including food products such as meat (Yayan et al., 2015). Infection caused by P. aeruginosa, exceptionally resistant isolates seems challenging to treat due to its resistance to a variety of antibiotics, and patients can die from pneumonia caused by Pseudomonas spp. (Tran et al., 2014). The increasing resistance of P.

aeruginosa to numerous antibiotics results from excessive antibiotic administration in both humans and animals, leading to the accumulation of antibiotic resistance and cross-resistance between antibiotics and the appearance of multidrug-resistant (MDR) forms of P. aeruginosa. This study's finding is in line with Atuanya et al. (2018), who reported resistant P.

aeruginosa isolated from slaughterhouse Effluents around Ikpoba River in Benin City, Nigeria. This is contrary to the findings of Jega et al. (2018), who reported the sensitivity of P. Aeruginosa isolated from slaughterhouse Effluent-Impacted Tagangu River, Aliero to both Chloramphenicol and Septrin. CONCLUSION The occurrence of antimicrobial-resistant pathogens such as E. coli, E. aerogenes, and P.

aeruginosa from this slaughterhouse points showed the risks associated with

antimicrobial **drug resistance transferred from** food- producing animals to humans. It is recommended that the management concern, such as local government health officers, **and community development officers** should increase on the sensitization of slaughterhouse workers by organizing conferences or conducting radio talk to educate these slaughterhouse workers on the treatment of slaughterhouse wastewater and health risk associated with antimicrobial **drug resistance transferred from** food-producing animals to humans.

ACKNOWLEDGMENT The author thanks all staff of the Microbiology laboratory of **Kebbi State University of Science and Technology**, Aliero, for their kind support during

INTERNET SOURCES:

- <1% - www.ncbi.nlm.nih.gov/pmc/articles/PMC3187576
- <1% - quizlet.com/101367472/chapter-15-methods-of...
- 2% - www.sciencedirect.com/science/article/pii/S...
- <1% - www.ijcmas.com/vol-3-9/David N Ogbonna.pdf
- <1% - www.nature.com/articles/s41598-020-65766-8
- <1% - www.britannica.com/science/latitude
- <1% - bdvets.binbd.com/JAVAR/V4I3/d215_pp227-233.pdf
- <1% - www.ncbi.nlm.nih.gov/pmc/articles/PMC3658817
- <1% - www.hindawi.com/journals/tswj/2012/764563/abs
- <1% - pgform.ksustapgs.net/reg/printprofile.php?studentidwiie...
- <1% - www.sciencedirect.com/.../coliform-bacteria
- <1% - quizlet.com/152140607/lab-4-standard-plate-count...
- <1% - onlinelibrary.wiley.com/doi/pdf/10.1111/jvim.13949
- <1% - fire.biol.wvu.edu/brodham/biol346_S07/labman_week9b.pdf
- <1% - www.ncbi.nlm.nih.gov/pmc/articles/PMC6787911
- 1% - academicjournals.org/journal/AJMR/article-full...
- <1% - www.ijser.org/researchpaper/Examination-of...
- <1% - www.ncbi.nlm.nih.gov/pmc/articles/PMC4948424
- <1% - quizlet.com/412837783/micro-lab-practical-2...
- <1% - quizlet.com/103092600/lab-quiz-2-flash-cards
- <1% - quizlet.com/129569452/micro-lab-exam-2-flash-cards
- 1% - this confirmatory **test was carried out to determine if the coliforms were of the fecal origin or** not.
- <1% - dramohammadi.ir/Download/IDFlowcharts.pdf
- <1% - www.ncbi.nlm.nih.gov/pmc/articles/PMC6449141
- <1% - pdfs.semanticscholar.org/b56c/857e8a19a857d0e98f...
- <1% - www.nature.com/articles/s41598-019-51907-1

1% - The Associated Press via Yahoo Sports
<1% - www.ncbi.nlm.nih.gov/pubmed/7006063
<1% - bmcrenotes.biomedcentral.com/articles/10.1186/s...
<1% - www.ncbi.nlm.nih.gov/pmc/articles/PMC6463849
<1% - www.ncbi.nlm.nih.gov/pubmed/31413825
<1% - quizlet.com/26940272/enterobacteriaceae-flash-cards
<1% - www.scirp.org/journal/PaperInformation.aspx?...
1% - journals.plos.org/plosone/article?id=10.1371/...
<1% - www.ncbi.nlm.nih.gov/pmc/articles/PMC3528471