

Chemical Constituents and Antioxidant Activity of *Melothria scabra* Naudin Fruits

Harni Sartika Kamaruddin ¹ Megawati ² Nurliana ¹Carla Wulandari Sabandar ^{1*} 

¹Department of Pharmacy, Universitas Sembilanbelas November Kolaka, Kolaka, Southeast Sulawesi, Indonesia

²Department of Chemistry, Universitas Sembilanbelas November Kolaka, Kolaka, Southeast Sulawesi, Indonesia

*email: carla@usn.ac.id

Keywords:

Antioxidant activity
Cucurbitaceae
Cucurbitacin B glycoside
Melothria scabra Naudin

Abstract

The fruit of *Melothria scabra* Naudin is traditionally used by natives of South East Sulawesi and has economic values in the local markets. Nonetheless, little scientific information was gained from this plant to support its development for nutraceutical and pharmaceutical aspects. This study aimed to investigate the phytochemicals contained in the ethanol extract and organic fractions (methanol, ethyl acetate, and hexane) of the fruits using specific reagents and an LC-MS/MS analysis, as well as to evaluate their total phenolics, total flavonoids, and DPPH radical scavenging activity using a dot-blot staining and spectrophotometric assays. Results showed that the fruits of *M. scabra* contained alkaloids, tannins, flavonoids, terpenoids, and saponins. Six compounds were successfully identified from the ethanol extract of the fruits for the first time that is D-1-[(3-carboxypropyl)amino]-1-deoxyfructose (1), fructose-C₃H₅NO (2), valine (3), 1 β , 3 α , 9 β -trihydroxyeudesma-5,11(13)-dien-12-oic acid (4), cucurbitacin B-2-O- α -L-rhamnopyranosyl- β -D-glucopyranoside (5), and 2-heptyl-3-hydroxy-4(H)-quinolone (6). Total phenolics in the extract and organic fractions were in the range of 54.2 \pm 2.4 to 259.1 \pm 8.4 mg GAE/g, while total flavonoids were in the range of 1.6 \pm 0.2 to 22.4 \pm 0.2 mg QE/g. The ethanol extract and its organic fractions (methanol and ethyl acetate) were potent radical scavengers with SC₅₀ values ranging from 20.7 to 37.5 μ g/mL when compared with ascorbic acid, gallic acid, and quercetin (SC₅₀ of 2.8 to 9.4 μ g/mL). This study concludes that *M. scabra* fruits could be developed as a source of natural antioxidant agents for nutraceuticals and pharmaceuticals purposes.

Received: October 6th, 2021

Accepted: November 13th, 2021

Published: November 30th, 2021



© 2021 Harni Sartika Kamaruddin, Megawati, Nurliana, Carla Wulandari Sabandar. Published by Institute for Research and Community Services Universitas Muhammadiyah Palangkaraya. This is an Open Access article under the CC-BY-SA License (<http://creativecommons.org/licenses/by-sa/4.0/>). DOI: <https://doi.org/10.33084/bjop.v4i4.2890>

INTRODUCTION

Melothria scabra Naudin belongs to the Cucurbitaceae family. It was originally from Mexico and Central America but has spread throughout Asia, including Indonesia. The plant also has some synonym names, which are *M. donnell-smithii* Cogn. Ex Donn.Sm., *M. donnell-smithii* var. *hirtella* Cogn., and *M. donnell-smithii* var. *rotundifolia* Cogn¹. The fruit of *M. scabra* is slightly bittersweet and has a unique aroma. Morphologically,

the fruit is oval with 3.5 in length and 2.5 cm in wide. It resembles a miniature version of watermelon and is locally known as “balongga” (Kolaka and Konawe) or “balongkha” (Buton). The fruit is consumed directly or processed as pickles and vegetables by natives of Southeast Sulawesi; hence, it is often found in local markets². In Sulawesi medicinal folklore, it is used to reduce blood pressure, while in Pakistan, it is applied as a mosquito repellent³. A previous study has reported the phytochemical screening and anti-diabetes activity of the

ethanol extract of its leaves⁴. In the present study, we reported the phytochemical constituents of the ethanol extract of the fruits analyzed using LC-MS/MS spectrometry and evaluated the antioxidant activity and organic fractions using total phenolics, total flavonoids, and 2,2-diphenyl-1-picrylhydrazyl radical assays. The study of the fruits of *M. scabra* is reported for the first time in this paper.

MATERIALS AND METHODS

Materials

Chemicals

Methanol, ethyl acetate, and *n*-hexane were technical grades and distilled before being used. Ethanol 96% was purchased locally. Folin-Ciocalteu reagent, aluminum chloride, gallic acid, quercetin, ascorbic acid, dimethyl sulfoxide (DMSO), and thin-layer chromatography (TLC) plate (silica gel 60 GF254, 0.25 mm) were purchased from Merck (Darmstadt, Germany). 2,2-diphenyl-1-picrylhydrazyl (DPPH) was obtained from HiMedia (Mumbai, India).

Plant material

Fresh fruits of *M. scabra* (3.329 kg) were collected from Tanggetada District of Kolaka Regency, Southeast Sulawesi, Indonesia, in July 2021. The morphology of fruits, leaves, flowers, and seedlings of *M. scabra* was presented in **Figure 1**. The fruit sample was identified, and its specimen was deposited at the Laboratorium Terpadu USN Kolaka with voucher number MS001. The collected fruits were washed with tap water and drained. Cleaned fruits were slidely cut and dried using the oven at 40°C for four days. Then, the dried samples were coarsely ground using a blender, weighted, and stored in an airtight plastic sealer until used.




Figure 1. Fruits (A), leaves (B), flower (C), and seedling (D) of *M. scabra*

Methods

Extraction and fractionation

The coarsely powdered fruits of *M. scabra* (295 g) were macerated using ethanol 96% for 3 x 48 hours at room temperature. The liquid extract was filtered using Whatman filter paper and concentrated using a vacuum rotary evaporator, yielding a yellowish-brown crude ethanol extract. For fractionation using three different polarities of solvents (methanol, ethyl acetate, *n*-hexane), 10 g of crude ethanol extract was firstly reconstituted in methanol (250 mL) and successively partitioned with *n*-hexane and ethyl acetate each for 3 x 250 mL in a separating funnel. All solvents were then evaporated to yield organic fractions of methanol, ethyl acetate, and *n*-hexane. Both extract and organic fractions were stored in amber bottles and kept in the refrigerator at 4°C until used.

Phytochemical screening

The phytochemicals contained in ethanol extract and organic fractions of *M. scabra* were screened according to our previous method⁵. The screening was carried out to detect the presence of alkaloids, tannins, flavonoids, terpenoids, steroids, and saponins.

LC-MS/MS analysis

The LC-MS/MS measurement of the ethanol extract of *M. scabra* was in collaboration with the Indonesian Institute of Sciences (LIPI). The method was adopted from our previous study with modifications⁶. The chemical constituents in the ethanol extract of *M. scabra* were separated using a reverse-phase column (ACQUITY UPLC BEH C8, 1.7 μm \times 100 mm) and gradiently eluted with water containing 0.1% formic acid (A) and acetonitrile containing 0.1% formic acid (B) at a flow rate of 0.3 mL/minute and column temperature of 40.0°C. The elution time was running for 16 minutes as follows: 95% A (0–1 minutes); 60% A (8 minutes); 100% B (11–13 minutes); and 5% B (16 minutes). The separated peaks were detected using a Xevo G2-XS QToF mass spectrometer. The mass experiments were carried out using a positive mode electrospray ionization (ESI) with acquisition parameters as follows: capillary voltage, 2 kV; cone voltage, 30 V; source temperature, 120°C; desolvation temperature, 500°C; cone gas flow, 50 L/h; desolvation gas flow, 1000 L/h. Experiments were carried out in two functions of MSE. Function one was carried out at a high CE ramp of 10 to 40 eV, cone voltage of 30 V, and collision energy of 6 eV. Meanwhile, function two was carried out at a high CE ramp of 10 to 40 eV, cone voltage of 30 V, and collision energy of 10 to 40 eV. For both functions, the acquisition time ran for 16 minutes with detection of mass-to-charge ratio (m/z) starting from m/z 50 to m/z 1200. The obtained m/z values of separated peaks were identified using the UNIFI software and analyzed using online mass databases, including the MetFrag, METLIN, PubChem, mzCloud dan ChemSpider.

Total phenolics assay

According to our previous study with modifications^{5,7}, total phenolics in the ethanol extract and organic fractions of *M. scabra* fruits were evaluated spectrophotometrically

using the Folin-Ciocalteu (FC) method. A linear gallic acid standard curve was first established within a concentration range of 6.25 to 100 $\mu\text{g}/\text{mL}$ versus absorbance of molybdenum-tungsten blue complex at 635 nm. **Equation 1** obtained as follows:

$$Abs = \left(0.0511 \times [\text{gallic acid in } \frac{\mu\text{g}}{\text{mL}}]\right) + 0.1036 \quad (r^2 = 0.990) \quad [1]$$

Stock solutions of samples (10 mg/mL) were prepared in DMSO, and the final concentration in the reaction mixture was 1 mg/mL. A volume of 0.2 mL of sample solution was triplicately transferred into vials and incubated with 0.2 mL of FC reagent for 5 minutes at room temperature. The mixture was then added with 0.8 mL NaHCO_3 solution (7.5%, w/v) and re-incubated for 30 minutes in the dark at room temperature. The absorbance was read against a sample blank containing sample solution and water. The total phenolics in the sample were obtained by the interpolation of absorbance in the equation and expressed as mg of gallic acid equivalent for each gram of sample (mg GAE/g).

Total flavonoids assay

Total flavonoids content in the ethanol extract and organic fractions of *M. scabra* fruits was evaluated spectrophotometrically using the aluminum chloride method as previously reported with modifications^{5,7}. A linear quercetin standard curve was first established within a concentration range of 1.56 to 100 $\mu\text{g}/\text{mL}$ versus the absorbance of the yellow complex at 425 nm. **Equation 2** obtained as follows:

$$Abs = \left(0.0799 \times [\text{quercetin in } \frac{\mu\text{g}}{\text{mL}}]\right) - 0.0688 \quad (r^2 = 0.999) \quad [2]$$

Stock solutions of samples (10 mg/mL) were prepared in DMSO, and the final concentration in the reaction mixture was 1 mg/mL. A volume of 1 mL of sample solution was triplicately transferred into vials and incubated with 1 mL of AlCl_3 solution (2%, w/v) for 15 minutes at room temperature. The absorbance was read against a sample blank containing sample solution and

water. The total flavonoids in the sample were obtained by the interpolation of absorbance in the equation and expressed as mg of quercetin equivalent for each gram of sample (mg QE/g).

DPPH assay

DPPH assay was carried out to determine the radical scavenging activity of ethanol extract and organic fractions of *M. scabra* fruits towards free radicals⁵⁷. The assay was performed qualitatively using dot-blot staining on TLC plate and quantitatively using the spectrophotometric method. For the qualitative assay, samples solutions (2 mg/mL) were prepared in their soluble solvents and spotted (20 µL) on a TLC plate with a final concentrations range of 6.2 to 100 µg/spot. After drying the solvents, the TLC plate was dipped into DPPH methanolic solution (0.5 mM) for 10 seconds, shortly dried using an air-dryer, and incubated for 30 minutes in the dark at room temperature. The radical scavenging activity was analyzed by observing the clear zone around the spot under daylight against the purple background of DPPH.

For quantitative analysis, samples stock solutions (10 mg/mL) were prepared in DMSO and serially diluted into eight concentrations ranging from 100 to 0.8 µg/mL. Triplicate volumes (1 mL) of diluted solutions were transferred into vials and incubated, each with 1 mL of DPPH solution (0.25 mM) for 15 minutes in the dark at room temperature. The absorbance was recorded at 515 nm against samples and blank containing DMSO. Sample absorbance (Abs_{sample}) was obtained after blank sample correction. The 50% radical scavenging concentration (SC_{50}) values of ethanol extract, organic fractions, and positive controls (ascorbic acid, gallic acid, quercetin) were calculated using GraphPad Prism 5 software (GraphPad Inc., California, US). The radical scavenging activity (%RSA) was calculated using **Equation 3** as follows:

$$\%RSA = \left(\frac{Abs_{DPPH\ control} - Abs_{Sample}}{Abs_{DPPH\ control}} \right) \times 100 \quad [3]$$

Statistical analysis

Data for all assays were presented as mean ± standard deviation (SD), which was obtained from three replications. Data were statistically analyzed using the GraphPad Prism 5 Software. Differences among groups were evaluated using Analysis of Variance (ANOVA) with Tukey's Test ($p < 0.05$). The significances among parameters (total phenolics, total flavonoids, DPPH) were measured using the Pearson correlation⁸.

RESULTS AND DISCUSSION

Extraction yield and phytochemical screening

The extraction of fruits of *M. scabra* was done by using ethanol 96% using the maceration technique. Aqueous ethanol was considered a less toxic solvent for further *in vitro* and *in vivo* studies based on cells and animal models⁹. The extract was obtained as a yellowish-brown gum with a percentage yield of 16.8%, showing a high number of polar compounds. This amount was observed in the yield of methanol fraction with a percentage yield of 67.9% compared to the ethyl acetate (27.0%) and hexane (5.0%) fractions, as presented in **Table I**.

Fruits of *M. scabra* also showed the presence of alkaloids, tannins, flavonoids, terpenoids, and saponins. Meanwhile, steroids were not detected, probably due to their low amounts. Distributions of all present groups of compounds varied among organic fractions, but alkaloids, terpenoids, and saponins seem dominants. Flavonoids represented as antioxidants¹⁰ were detected in polar and semipolar solvents extraction, including ethanol extract, methanol, and ethyl acetate fractions. The occurrence of alkaloids and flavonoids has been reported from the leaves of this plant, along with glycosides, carbohydrates, and proteins⁴.


Table I. Phytochemical groups detected in the ethanol extracts and organic fractions of *M. scabra* fruits

Samples	Yield (g)	Phytochemical groups
Ethanol extract	49.53	Alkaloids, tannins, flavonoids, terpenoids, saponins
Metanol fraction	6.79	Alkaloids, tannins, flavonoids, terpenoids, saponins
Ethyl acetate fraction	2.70	Alkaloids, flavonoids, terpenoids, saponins
<i>n</i> -hexane fraction	0.50	Alkaloids, terpenoids, saponins

Identification of chemical constituents

The chromatographic separation of chemical constituents in the ethanol extract of *M. scabra* fruits was analyzed using the ultra-performance liquid chromatography (UPLC) method, which separated various peaks starting from retention time (Rt) of 1.5 to 12.5 minutes. Some major peaks were observed at the initial retention time from 1.5 to 2.5 minutes, indicating the composition of polar compounds. The UPLC chromatogram of these peaks and other minors is displayed in **Figure 2**.

Further detection using a mass detector yielded the separated peaks' mass-to-charge ratio (m/z). As many as six compounds (**1-6**) were successfully identified from the ethanol extract of *M. scabra* fruit, and most were polar compounds (**1-5**). The LC-MS/MS identification of these compounds was displayed in **Table II**, and their two-dimensional structures were illustrated in **Figure 3**.

**Figure 2.** UPLC chromatogram of separated peaks of compounds in the ethanol extract of *M. scabra* fruits

Two major peaks were observed at the retention time of 1.7 and 1.80 minutes, having protonated pseudo-molecular ion of m/z 266 and m/z 1.80. The fragmentation of these ions resulted in the identification of compounds **1** and **2**, classified as Heyns compounds containing reducing sugar (fructose) and amino acid¹¹.

As far as our concern, both compounds have yet previously reported from plants of the Cucurbitaceae family but found as the main components in the jujube (Rhamnaceae) and mulberry (Moraceae) fruits analyzed using a high-sensitive HPLC-ESI-Q-ToF-MS/MS¹². Compound **1** was also found in the cured tobacco leaves (Solanaceae) as well as stored apricots and peaches fruits (Rosaceae)¹³. Meanwhile, compound **2** was predicted as a product of the Maillard reaction of fructose with primary or secondary amines, in which its exact structure needs further identification¹².

Table II. Chemical constituents of the ethanol extract of *M. scabra* fruits identified using LC-MS/MS

Rt (min)	(+)-ESI	Identified constituents	MS/MS fragmentation (m/z)
1.73	266.12362 [M+H] ⁺	D-1-[(3-Carboxypropyl)amino]-1-deoxyfructose (1) C ₁₀ H ₁₉ NO ₇ MW 265.2604	248/206/104
1.80	234.09724 [M+H] ⁺	Fructose-C ₃ H ₅ NO (2) C ₉ H ₁₅ NO ₆ MW 233.2185	216/159
1.85	118.08588 [M+H] ⁺	Valine (3) C ₃ H ₁₁ NO ₂ MW 117.14634	102/100/86/84
1.93	305.13435 [M+Na] ⁺	1 β ,3 α ,9 β -Trihydroxyeudesma-5,11(13)-dien-12-oic acid (4) C ₁₅ H ₂₂ O ₅ MW 282.3322	305/227/191
7.85	865.42170 [M] ⁺	Cucurbitacin B-2-O- α -L-rhamnopyranosyl- β -D-glucopyranoside (5) C ₄₄ H ₆₆ O ₁₇ MW 866.9846	663/501/483/465/ 441/336/177
10.27	260.16444 [M+H] ⁺	2-Heptyl-3-hydroxy-4(H)-quinolone (6) C ₁₆ H ₂₁ NO ₂ MW 259.3434	173/146

The fruits of *M. scabra* also produced a significant count of an essential amino acid valine (**3**), observed as one of the major peaks at a retention time of 1.85 minutes and had an m/z 118 [M+H]⁺. Hence, the fruits of *M. scabra* could be developed as a source of nutritive essential amino acids. Another major peak was observed at a retention time of 1.93 minutes, with an adducted ion m/z

305 [M+Na]⁺. The compound was identified as trihydroxyeudesma-5,11(13)-dien-12-oic acid (4) and classified as a sesquiterpenoid. The presence of this compound might be attributed to a sweet and unique aroma of the fruits of *M. scabra*.




Figure 3. Chemical compounds identified in the ethanol extract of *M. scabra* fruits

The occurrence of a saponin (5) was detected at a retention time of 7.85 minutes and had an m/z 865 [M]⁺. The fragmentation of this compound yielded fragment ions at m/z 663, 603, and 501 (Figure 4), which correspond to the successive eliminations of rhamnosyl and acetyl groups and fission of a glucose ring, following its loss (Figure 5). The loss of the acetyl group (-C₂H₅O) suggested the main skeleton of this compound as cucurbitacin B glycoside^{14,15}, which is further confirmed by fragment ions at m/z 483 and 465 due to the loss of two hydroxyl groups. The cucurbitacins (cucurbitane-type) are highly oxygenated tetracyclic triterpenoids with a wide range of promising biological activities, including cytotoxicity, hepatoprotective, antidiabetic effects, cardiovascular protection, antioxidant, and anti-

inflammatory¹⁵. This group of compounds is considered as chemical markers for the Cucurbitaceae family¹⁶.




Figure 4. LC-MS/MS spectrum of compound 5




Figure 5. The plausible MS² fragmentation of compound 5

Compound **5** was further identified as cucurbitacin B-2-O- α -L-rhamnopyranosyl- β -D-glucopyranoside. The occurrence of cucurbitacin B-2-O- β -D-glucopyranoside has been reported from the leaves of *Citrullus colocynthis* and showed the inhibitory effect of the growth of human breast cancer cells¹⁷. However, the occurrence of compound **5** as cucurbitacin B diglycoside has not been previously reported. Nonetheless, its exact structure needs further studies by isolation and identification using more spectroscopic measurements.

One non-polar peak at a retention time of 10.27 minutes had an m/z value of 260 [M+H]⁺ as the base peak. The peak showed fragment ions at m/z 173 and 146, corresponding to the loss of heptyl and hydroxyl side groups from the quinolone skeleton. This peak was identified as 2-heptyl-3-hydroxy-4(H)-quinolone (**6**) and belonged to the quinolone alkaloid group. The occurrence of compound **6** in the fruits of *M. scabra* is considered unique since the compound has been acknowledged as a quorum sensing (QS) produced by plant growth-promoting bacteria such as *Pseudomonas* and *Paenibacillus* species. We suggested compound **6** as a diffusible signal in communication between *M. scabra* and soil bacteria, resulting in the protection of the fruits from other pathogenic bacteria and fungi since they are growing on the soil's surface¹⁸. In addition, as a QS, quinolones were also produced by plants of the Rutaceae family which exhibited remarkable antimicrobial activity¹⁹. Hence, further studies on the antimicrobial activity of *M. scabra* fruits could be developed.

Total phenolics and total flavonoids

The fruits of *M. scabra* contained a high amount of phenolics, especially in the methanol and ethyl acetate fractions (Table III). Moreover, the ethyl acetate fraction contained more flavonoids than other fractions, indicating the accumulation of semi-polar flavonoids in the ethanol extract of *M. scabra* fruits²⁰. High contents of

phenolics and flavonoids might be contributed to the antioxidant activity¹⁰.

Table III. Total phenolics and total flavonoids in the ethanol extract and organic fractions of *M. scabra* fruits

Samples	Total phenolics (mg GAE/g, mean \pm SD, n=3)	Total flavonoids (mg GAE/g, mean \pm SD, n=3)
Ethanol extract	86.3 \pm 3.4	15.5 \pm 0.3
Methanol fraction	259.1 \pm 8.4	17.2 \pm 0.4
Ethyl acetate fraction	242.9 \pm 3.3	22.4 \pm 0.2
n-hexane fraction	54.2 \pm 2.4	1.6 \pm 0.2

DPPH radical scavenging activity

The ability of *M. scabra* fruits to scavenge DPPH radicals was evaluated using qualitative and quantitative methods. Qualitatively, potent scavenging activity against DPPH radical was shown by the ethanol extract, methanol, and ethyl acetate fractions compared to ascorbic acid as the positive control (Figure 6). Meanwhile, the hexane fraction showed weak antioxidant activity. Their activities were concentration-dependent at a concentration range of 100 to 6.2 μ g/spot.




Figure 6. DPPH dot-blot staining of ethanol extract and organic fractions of *M. scabra* fruit

In line with the qualitative assay, the ethanol extract and organic fractions (methanol and ethyl acetate) of *M. scabra* fruits quantitatively exhibited potent antioxidant activity compared to the hexane fraction (Table IV). The ethyl acetate fraction was considered the most potent radical

scavenger with an SC_{50} value of 20.7 $\mu\text{g}/\text{mL}$, followed by the methanol fraction (SC_{50} 28.6 $\mu\text{g}/\text{mL}$) and the ethanol extract (SC_{50} 37.5 $\mu\text{g}/\text{mL}$). As the positive controls, their activities were comparable to ascorbic acid, gallic acid, and quercetin.

Table IV. DPPH radical scavenging activity of the ethanol extract and organic fractions of *M. scabra* fruits

Samples	%RSA (at 100 $\mu\text{g}/\text{mL}$, mean \pm SD, n=3)	SC_{50} ($\mu\text{g}/\text{mL}$)
Ethanol extract	62.7 \pm 0.6 [*]	37.5
Methanol fraction	67.6 \pm 0.9 [*]	28.6
Ethyl acetate fraction	80.5 \pm 2.0 [*]	20.7
<i>n</i> -hexane fraction	49.1 \pm 0.2 [*]	476.3 [*]
Ascorbic acid	98.8 \pm 0.6	2.8
Gallic acid	97.2 \pm 0.3	3.9
Quercetin	98.1 \pm 0.6	9.4

^{*}Values are significantly different compared to positive controls ($p < 0.05$) based on Tukey's test.

The correlations of phenolics and flavonoids in the fruits of *M. scabra* at high concentration (100 $\mu\text{g}/\text{mL}$) were considered moderate analyzed using the Pearson correlation with r values of 0.8274 ($p = 0.1726$) and 0.8226 ($p = 0.1774$), respectively (Table V). However, these contents found not to correlate with the SC_{50} values ($r = -0.6920$, $p = 0.3080$ for phenolics and $r = -0.3695$, $p = 0.6305$ for flavonoids). Results indicated that phenolics and flavonoids did not solely determine the radical scavenging activity of the ethanol extract and its organic fraction. Other compounds may also contribute to the activity, either scavenging the radical or assisting the radical activity²¹. Hence, further compounds isolation and structure elucidation as well as assessments of antioxidant activity using other assays, especially *in vitro* cell-based assay and *in vivo* assay using animal models, need to be carried out.

Table V. Pearson correlation (r) and linear regression curve (r^2) between total phenolics and total flavonoids and DPPH ($p < 0.05$, two-tailed)

Content	%RSA (100 $\mu\text{g}/\text{mL}$)		SC_{50}	
	Pearson (r)	Linear curve (r^2)	Pearson (r)	Linear curve (r^2)
Total phenolics	0.8274	0.6847	-0.6920	0.4789
Total flavonoids	0.8226	0.6766	-0.3695	0.1365

CONCLUSION

This study concludes that the fruits of *M. scabra* contained alkaloids, tannins, flavonoids, terpenoids, and saponins. Among these, six compounds were identified as D-1-[(3-carboxypropyl)amino]-1-deoxyfructose (1), fructose- C_3H_5NO (2), valine (3), 1 β , 3 α , 9 β -trihydroxyeudesma-5,11(13)-dien-12-oic acid (4), cucurbitacin B-2-O- α -L-rhamnopyranosyl- β -D-glucopyranoside (5), and 2-heptyl-3-hydroxy-4(H)-quinolone (6). These compounds in *M. scabra* are reported for the first time in this study. The fruits also contained a high amount of phenolics and flavonoids, which moderately correlated to their radical scavenging activity. The ethanol extract, methanol, and ethyl acetate fractions showed potent antioxidant activity, which could be developed as a source of natural antioxidant agents.

ACKNOWLEDGMENT

The authors would like to thank the Ministry of Education, Culture, Research, and Technology of the Republic of Indonesia and Badan Riset dan Inovasi Nasional (BRIN) for a research grant scheme (Penelitian Dosen Pemula 2021, Contract Number: 445/UN56D/PN.01.00/2021) for financial support.

AUTHORS' CONTRIBUTION

Harni Sartika Kamaruddin: the project leader responsible for research design, results validation, and editing the manuscript. **Megawati:** project team member who contributed ideas for research team and specifically provided technical assistance for extraction, fractionation, and chemical assessment. **Nurliana:** an undergraduate candidate who conducted the experimental work and data collection. **Carla Wulandari Sabandar:** project team member, contributed ideas for research design, conducted data analysis, and revised the manuscript.

DATA AVAILABILITY

The supporting data of the article are accessible from the corresponding author upon reasonable request.

CONFLICT OF INTEREST

The author declares that there is no conflict of interest.

The author is fully responsible for the content and writing of this article.

REFERENCES

- World Flora Online. *Melothria scabra* Naudin [Internet]; 2021 [updated 2021; cited 2021 Oct 6]. Available from: <http://www.worldfloraonline.org/taxon/wfo-0000377170>
- Arniawati, Kahirun, Arafah N, Iryan. Kontribusi kawasan Taman Nasional Rawa Aopa Watumohai sebagai Penyedia Pangan bagi Masyarakat Desa Tatangge. *J Eco Green*. 2019;5(1):83-8.
- Ahmed I, Nasir S, Yousaf I, Ahmad B, Zafar S, Javaid A. Plant Extracts along with Selective Chemicals and *Bacillus thuringiensis israelensis*: A Novel Approach to Tackle the Problem of Insecticidal Resistance in Mosquitoes. *Pak J Agri Sci*. 2019;56(4):905-11. doi:10.21162/PAKJAS/19.8481
- Anusha G, Sunayana R, Ponnampalani M, Kumar BA. Phytochemical Investigation and In Vitro Antidiabetic Activity of *Melothria scabra*. *Asian J Pharm Res Dev*. 2019;7(4):43-48. doi:10.22270/ajprd.v7i4.553
- Sabandar CW, Jalil J, Ahmat N, Aladdin N-A, Kamaruddin S, Wahyuningrum R. Aktivitas Antioxidant dan Penghambatan Xantin Oksidase Kulit Batang Songi (*Dillenia serrata* Thunb.). *J Farmasi Galenika*. 2020;6(1):151-9. doi:10.22487/j24428744.2020.v6.i1.15008
- Sahidin I, Sabandar CW, Wahyuni, Hamsidi R, Mardikasari SA, Zubaydah WOS, et al. Investigation of Compounds and Biological Activity of Selected Indonesian Marine Sponges. *Nat Prod J*. 2019;10(3):312-21. doi:10.2174/2210315509666190627105237
- Sabandar CW, Jalil J, Ahmat N, Aladdin N-A. Assessment of Antioxidant and Xanthine Oxidase Inhibitory Activity of *Triadia conchinchinensis* Stem Bark. *Curr Res Biosci Biotechnol*. 2019;1(1):39-44.
- Al-Farsi M, Al-Amri A, Al-Hadhrami A, Al-Belushi S. Color, flavonoids, phenolics and antioxidants of Omani honey. *Heliyon*. 2018;4(10):e00874. doi:10.1016/j.heliyon.2018.e00874
- Kraut JA, Mullins ME. Toxic Alcohols. *N Engl J Med*. 2018;378(3):270-80. doi:10.1056/nejmra1615295
- Heim KE, Tagliaferro AR, Bobilya DJ. Flavonoid Antioxidants: Chemistry, Metabolism and Structure-Activity Relationships. *J Nutr Biochem*. 2002;13(10):572-84. doi:10.1016/s0955-2863(02)00208-5
- Yuan H, Sun L, Chen M, Wang J. The Simultaneous Analysis of Amadori and Heyns Compounds in Dried Fruits by High Performance Liquid Chromatography Tandem Mass Spectrometry. *Food Anal Methods*. 2017;10:1097-105. doi:10.1007/s12161-016-0669-1
- Li R, Wei M, Guo G, Li Y, Pan X, Song X, et al. Analysis of Main Components in Jujube and Mulberry Extracts by High-Sensitive HPLC-ESI-Q-TOF-MS/MS. *J Chromatogr Sci*. 2021;59(9):806-12. doi:10.1093/chromsci/bmaa133
- Collins PM. Dictionary of Carbohydrates with CD-ROM. 2nd edition. Boca Raton, Florida: Chapman & Hall; 2006.
- Lang KL, Silva IT, Zimmermann LA, Machado VR, Teixeira MR, Lapuh MI, et al. Synthesis and Cytotoxic Activity Evaluation of Dihydrocucurbitacin B and Cucurbitacin B Derivatives. *Bioorg Med Chem*. 2012;20(9):3016-30. doi:10.1016/j.bmc.2012.03.001
- Tannin-Spitz T, Bergman M, Grossman S. Cucurbitacin Glucosides: Antioxidant and Free-Radical Scavenging Activities. *Biochem Biophys Res Commun*. 2007;364(1):181-6. doi:10.1016/j.bbrc.2007.09.075
- Shah SSA, Hussain MI, Aslam MK, Rivera G. Natural Products; Pharmacological Importance of Family Cucurbitaceae: A Brief Review. *Mini Rev Med Chem*. 2014;14(8):694-705. doi:10.2174/1389557514666140820113055
- Tannin-Spitz T, Grossman S, Dovrat S, Gottlieb HE, Bergman M. Growth Inhibitory Activity of

- Cucurbitacin Glucosides Isolated from *Citrullus colocynthis* on Human Breast Cancer Cells. *Biochem Pharmacol.* 2007;73(1):56-67. doi:[10.1016/j.bcp.2006.09.012](https://doi.org/10.1016/j.bcp.2006.09.012)
18. Pierson LS, Pierson EA. Roles of Diffusible Signals in Communication among Plant-Associated Bacteria. *Phytopathology.* 2007;97(2):227-32. doi:[10.1094/PHYTO-97-2-0227](https://doi.org/10.1094/PHYTO-97-2-0227)
19. Ulubelen A, Öztürk M. Alkaloids and Coumarins from *Ruta* Species. *Nat Prod Commun.* 2006;1(10):851-7. doi:[10.1177/1934578X0600101006](https://doi.org/10.1177/1934578X0600101006)
20. Darmadi J, Batubara RR, Himawan S, Azizah NN, Audah HK, Arsianti A, et al. Evaluation of Indonesian mangrove *Xylocarpus granatum* leaves ethyl acetate extract as potential anticancer drug. *Sci Rep.* 2021;11:6080. doi:[10.1038/s41598-021-85383-3](https://doi.org/10.1038/s41598-021-85383-3)
21. Yang H, Dong Y, Du H, Shi H, Peng Y, Li X. Antioxidant Compounds from Propolis Collected in Anhui, China. *Molecules.* 2011;16(4):3444-55. doi:[10.3390/molecules16043444](https://doi.org/10.3390/molecules16043444)